

### MY NEXT STEPS

Provider appointment (if applies):

Da	te: Time:
Pro	ovider:
Pla	ace:
	one #:
How w	ill I get there? (circle)
a)	Drive myself
b)	Have a friend or family member drive me
c)	Public transportation
d)	Other:
What s	upport might I need to get to my
appoin	tment? (circle)
a)	Transportation
b)	Childcare
c)	Work
d)	Other:
What it	f I can't make my appointment?

#### **RESOURCES**

- Find a local health provider in your area: https://opa-fpclinicdb.hhs.gov/
- Find a health center: https://ncchca.site-ym.com/page/FindCHC
- NC Free Clinics: http://ncafcc.org/
- For more information about birth control methods (including emergency contraception) and other sexual health topics, visit: www.bedsider.org or www.fpntc.org
- For more information about Preconception Health visit: https://everywomannc.org/ or call 919-781-2481
- Department of Health and Human Services Customer Service Center can assist in finding programs and people to help you. Call 1-800-662-7030



www.ncdhhs.gov NCDHHS is an equal opportunity employer and provider. 02/2019

The creation of this material was supported, in part, by Title X funding. Supported by the NC Division of MH/DD/SAS through funding from SAMHSA State Targeted Response to the Opioid Crisis grant funds.

Adapted from D. Rinehart et al., NIDA R34DA039381
Developing and testing the feasibility, accessibility and initial efficacy of a brief peer led intervention to improve reproductive health among women in opioid medication-assisted treatment-"SHINE" study.



# Family Planning:

# Deciding If or When To Have Children







### **MY PLAN**

Today's Date:
Do you already have a child or children? ☐ Yes ☐ No
If yes, did you/partner give birth in the last year? □ Yes □ No
Do you want a child in the next year?
<ul><li>Yes. Talk to your healthcare provider about preparing for a healthy pregnancy.</li></ul>
<ul><li>No. Talk about Family Planning options.</li></ul>
☐ I don't know. Talk about Family Planning options and preparing for a healthy pregnancy.

Method:	
What I Like About It	What I'd Like to Know
a. Effectiveness	a. Effectiveness
b. Few potential	b. Potential side
side effects	effects
c. Cost	c. Cost
d. Other	d. Other

Method:	
What I Like About It	What I'd Like to Know
a. Effectiveness	a. Effectiveness
b. Few potential	b. Potential side
side effects	effects
c. Cost	c. Cost
d. Other	d. Other

Method:	
What I Like About It	What I'd Like to Know
a. Effectiveness	a. Effectiveness
b. Few potential	b. Potential side
side effects	effects
c. Cost	c. Cost
d. Other	d. Other

## WOULD YOU LIKE TO BECOME PREGNANT IN THE NEXT YEAR?

<b>+</b>	FAMILY PLANNING	METHOD	HOW LONG IT LASTS	I WOULD NEED TO DO THE FOLLOWING	I WOULD NEED TO THINK ABOUT	RISK OF PREGNANCY
NEVER	Sterilization (female or male)		Permanent	See my provider	If I want to get pregnant in the future, this is not a good method	Less than 1 in 100
	Abstinence* (female and male)			Not have sex right now	Having a lot of self-control in order not to become pregnant	0 in 100
	IUD (hormonal or hor Placed into uteru Can be taken ou anytime (female	ıs one	3-10 years	See my provider	Hormonal: Sometimes causes irregular bleeding or no bleeding     Hormone-free: Sometimes makes periods heavier and increases cramping	Less than 1 in 100
	Implant • Placed into arm • Can be taken out anytime (fen		1-3 years	See my provider     Do nothing until removed or replaced	Sometimes causes irregular bleeding or no bleeding	Less than 1 in 100
NOT IN THE NEXT YEAR	The Shot (female)	묩	3 months	See my provider every 3 months	Sometimes causes changes in bleeding and/or increases feeling hungry	6 in 100
	Ring (female)		1 month	Insert a ring into my vagina every month	Sometimes causes headaches, breast tenderness, nausea, or increase risk of blood clots     Must keep ring in a cool environment	9 in 100
IN THE	Patch (female)		1 week	Place a patch on my skin every week	Sometimes causes headaches, breast tenderness, nausea, or increase risk of blood clots	9 in 100
NOT	Pill (female)	000 000 000 000 000 000	1 day	Take a pill every day at the same time	Sometimes causes headaches, breast tenderness, nausea, or increase risk of blood clots.     Not a good method if you are 35 or older and use tobacco products	9 in 100
	<b>Diaphragm</b> (female)			Use with spermicide EVERY time I have sex	Must be used correctly     EVERY time you have sex     to be effective	12 in 100
	Condom* (male or female)		1 time	Use EVERY time     I have sex	Must be used correctly EVERY time you have sex to be effective	18 in 100
	Pulling out* (male)			Male withdraws before ejaculation	Female partners have no control over pulling out	22 in 100
	Rhythm method/ Fertility awareness*			Track my fertile days each month	May be difficult to determine the fertile time of the month	24 in 100

\*How will I talk with my partner(s) about this method?

Condom & abstinence are the only methods that protect against STDs.