

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Alamance County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	58.52	59.63	63.21	8.01	1,127	1,783	63.21	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.06	29.40	27.36	-5.85	417	1,524	27.36	28.75
B. Excessive Weight Gain	48.31	49.97	53.28	10.29	812	1,524	53.28	48.42
C. Inadequate Weight Gain	19.57	19.02	15.55	-20.54	237	1,524	15.55	19.51
D. Unknown Weight Gain	3.06	1.62	3.81	24.54	58	1,524	3.81	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.37	12.00	11.44	-30.10	204	1,783	11.44	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	77.58	75.78	79.49	2.46	735	941	78.11	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	33.71				359	1,065	33.71	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	10.20		0.57	-94.41	20	1,142	1.75	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Albemarle Regional Health Services District

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	53.21	65.82	65.98	24.00	923	1,399	65.98	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	20.49	27.48	28.12	37.24	338	1,202	28.12	28.75
B. Excessive Weight Gain	44.57	49.63	50.67	13.67	609	1,202	50.67	48.42
C. Inadequate Weight Gain	14.69	20.34	20.38	38.72	245	1,202	20.38	19.51
D. Unknown Weight Gain	20.24	2.54	0.83	-95.89	10	1,202	0.83	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.43	13.05	10.01	-4.05	140	1,399	10.01	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.31	68.28	82.29	15.41	664	794	83.63	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	23.35				187	801	23.35	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	7.07		7.20	1.80	79	820	9.63	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Bertie

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	67.74	72.56	73.37	8.31	124	169	73.37	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.39	20.15	22.46	-20.87	31	138	22.46	28.75
B. Excessive Weight Gain	44.52	50.00	50.00	12.32	69	138	50.00	48.42
C. Inadequate Weight Gain	26.45	29.10	27.54	4.10	38	138	27.54	19.51
D. Unknown Weight Gain	0.65	0.75	0.00	-100.00	0	138	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	7.53	12.80	10.06	33.64	17	169	10.06	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	80.26	80.81	88.01	9.66	115	132	87.12	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	9.59		5.80	-39.49	8	130	6.15	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Camden

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	47.37	69.39	66.67	40.74	60	90	66.67	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	22.22	29.03	41.56	87.01	32	77	41.56	28.75
B. Excessive Weight Gain	39.51	49.46	45.45	15.06	35	77	45.45	48.42
C. Inadequate Weight Gain	11.11	18.28	10.39	-6.49	8	77	10.39	19.51
D. Unknown Weight Gain	27.16	3.23	2.60	-90.44	2	77	2.60	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	9.47	11.22	12.22	29.01	11	90	12.22	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	85.39	71.43	72.28	-15.36	19	31	61.29	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			4.55		5	31	16.13	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Chowan

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.87	63.71	68.67	9.23	114	166	68.67	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	23.70	26.32	31.94	34.77	46	144	31.94	28.75
B. Excessive Weight Gain	54.81	54.39	39.58	-27.79	57	144	39.58	48.42
C. Inadequate Weight Gain	18.52	19.30	28.47	53.75	41	144	28.47	19.51
D. Unknown Weight Gain	2.96	0.00	0.00	-100.00	0	144	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	14.97	17.74	10.84	-27.57	18	166	10.84	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.90	77.97	86.05	9.07	90	106	84.91	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			1.61		2	103	1.94	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Currituck

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	33.59	68.46	77.31	130.13	201	260	77.31	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	12.77	23.21	28.13	120.31	63	224	28.13	28.75
B. Excessive Weight Gain	27.23	47.32	56.25	106.54	126	224	56.25	48.42
C. Inadequate Weight Gain	6.81	19.64	14.29	109.82	32	224	14.29	19.51
D. Unknown Weight Gain	53.19	9.82	1.34	-97.48	3	224	1.34	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.94	15.00	10.00	-8.57	26	260	10.00	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.93	58.61	71.15	-3.76	80	108	74.07	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						101		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Gates

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	26.67	57.69	55.05	106.42	60	109	55.05	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	8.86	30.86	23.91	169.88	22	92	23.91	28.75
B. Excessive Weight Gain	21.52	54.32	55.43	157.61	51	92	55.43	48.42
C. Inadequate Weight Gain	7.59	13.58	20.65	171.92	19	92	20.65	19.51
D. Unknown Weight Gain	62.03	1.23	0.00	-100.00	0	92	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.00	11.54	10.09	0.92	11	109	10.09	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.26	48.09	81.33	3.93	45	53	84.91	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						54		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Pasquotank

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	58.59	63.25	59.84	2.13	295	493	59.84	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	22.14	30.59	26.06	17.67	111	426	26.06	28.75
B. Excessive Weight Gain	53.57	48.59	52.35	-2.28	223	426	52.35	48.42
C. Inadequate Weight Gain	15.00	19.96	20.42	36.15	87	426	20.42	19.51
D. Unknown Weight Gain	9.29	0.87	1.17	-87.36	5	426	1.17	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	9.73	11.01	10.14	4.23	50	493	10.14	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	59.13	61.63	83.22	40.75	263	299	87.96	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	14.72		15.06	2.32	62	328	18.90	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Perquimans

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	60.56	68.55	61.61	1.72	69	112	61.61	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	22.50	29.46	32.67	45.21	33	101	32.67	28.75
B. Excessive Weight Gain	54.17	50.00	47.52	-12.26	48	101	47.52	48.42
C. Inadequate Weight Gain	16.67	20.54	19.80	18.81	20	101	19.80	19.51
D. Unknown Weight Gain	6.67	0.00	0.00	-100.00	0	101	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.27	16.13	6.25	-44.53	7	112	6.25	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.64	74.78	83.77	18.59	52	65	80.00	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.21		1.22	0.41	2	73	2.74	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Alexander County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	71.51	70.94	66.18	-7.45	227	343	66.18	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.25	28.19	30.26	7.14	92	304	30.26	28.75
B. Excessive Weight Gain	56.82	50.00	51.64	-9.11	157	304	51.64	48.42
C. Inadequate Weight Gain	13.31	21.14	17.11	28.50	52	304	17.11	19.51
D. Unknown Weight Gain	1.62	0.67	0.99	-39.21	3	304	0.99	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	19.83	19.09	18.95	-4.45	65	343	18.95	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.27	73.16	70.62	-3.61	150	211	71.09	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	40.47				87	215	40.47	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	0.95			-100.00		217		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Anson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	54.20	52.32	57.83	6.69	133	230	57.83	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.35	23.76	20.00	-26.89	40	200	20.00	28.75
B. Excessive Weight Gain	52.91	53.96	51.50	-2.67	103	200	51.50	48.42
C. Inadequate Weight Gain	19.28	22.28	28.00	45.21	56	200	28.00	19.51
D. Unknown Weight Gain	0.45	0.00	0.50	11.50	1	200	0.50	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.79	16.88	13.91	-17.15	32	230	13.91	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	83.14	84.25	82.36	-0.94	165	190	86.84	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	44.62				87	195	44.62	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	4.33			-100.00		204		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Appalachian District

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	78.02	77.60	78.01	-0.02	525	673	78.01	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.91	28.78	29.60	-4.25	169	571	29.60	28.75
B. Excessive Weight Gain	46.11	46.02	47.46	2.92	271	571	47.46	48.42
C. Inadequate Weight Gain	19.26	21.79	22.24	15.50	127	571	22.24	19.51
D. Unknown Weight Gain	3.72	3.41	0.70	-81.15	4	571	0.70	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.57	15.90	13.37	-1.45	90	673	13.37	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	64.42	73.61	75.96	17.92	267	341	78.30	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.34				98	372	26.34	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	6.68		3.11	-53.46	3	387	0.78	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Alleghany

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	54.26	72.73	69.23	27.60	72	104	69.23	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.97	31.03	31.40	20.87	27	86	31.40	28.75
B. Excessive Weight Gain	38.96	47.13	48.84	25.35	42	86	48.84	48.42
C. Inadequate Weight Gain	15.58	21.84	18.60	19.38	16	86	18.60	19.51
D. Unknown Weight Gain	19.48	0.00	1.16	-94.03	1	86	1.16	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.09	30.30	24.04	32.92	25	104	24.04	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	79.87	70.59	80.81	1.18	61	71	85.92	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	7.73		2.58	-66.68		80		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Ashe

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	80.41	73.54	78.04	-2.95	167	214	78.04	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.43	31.41	20.79	-33.86	37	178	20.79	28.75
B. Excessive Weight Gain	46.67	45.03	55.06	17.98	98	178	55.06	48.42
C. Inadequate Weight Gain	20.48	19.90	24.16	17.98	43	178	24.16	19.51
D. Unknown Weight Gain	1.43	3.66	0.00	-100.00	0	178	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	19.18	19.73	12.62	-34.23	27	214	12.62	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	74.89	74.79	75.06	0.22	93	123	75.61	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	8.24		3.95	-52.05	3	142	2.11	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Watauga

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	82.89	81.35	80.56	-2.81	286	355	80.56	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.80	26.71	34.20	7.54	105	307	34.20	28.75
B. Excessive Weight Gain	47.54	46.29	42.67	-10.24	131	307	42.67	48.42
C. Inadequate Weight Gain	19.34	22.85	22.15	14.50	68	307	22.15	19.51
D. Unknown Weight Gain	1.31	4.15	0.98	-25.49	3	307	0.98	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	8.26	9.73	10.70	29.60	38	355	10.70	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	49.05	73.52	74.73	52.36	113	147	76.87	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	4.81		2.55	-46.93		165		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Beaufort County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	76.84	70.02	74.07	-3.60	360	486	74.07	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.38	26.03	26.85	-1.92	105	391	26.85	28.75
B. Excessive Weight Gain	50.71	44.59	47.31	-6.70	185	391	47.31	48.42
C. Inadequate Weight Gain	17.38	26.29	25.32	45.67	99	391	25.32	19.51
D. Unknown Weight Gain	4.52	3.09	0.51	-88.69	2	391	0.51	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.91	17.00	16.67	29.10	81	486	16.67	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.76	82.58	86.16	9.41	271	298	90.94	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	27.71				87	314	27.71	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	21.09		34.88	65.44	166	342	48.54	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Bladen County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	59.23	50.69	56.41	-4.76	198	351	56.41	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.56	33.11	30.39	-0.58	86	283	30.39	28.75
B. Excessive Weight Gain	44.19	43.24	41.70	-5.64	118	283	41.70	48.42
C. Inadequate Weight Gain	23.59	22.64	27.56	16.85	78	283	27.56	19.51
D. Unknown Weight Gain	1.66	1.01	0.35	-78.73	1	283	0.35	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.73	15.98	19.66	4.94	69	351	19.66	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	83.05	83.68	78.65	-5.30	204	256	79.69	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.07				73	280	26.07	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.32		9.74	193.43	37	287	12.89	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Brunswick County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	65.27	63.43	56.23	-13.86	587	1,044	56.23	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.44	27.75	27.51	0.27	249	905	27.51	28.75
B. Excessive Weight Gain	51.32	53.28	51.82	0.98	469	905	51.82	48.42
C. Inadequate Weight Gain	20.09	18.11	20.22	0.64	183	905	20.22	19.51
D. Unknown Weight Gain	1.15	0.85	0.44	-61.50	4	905	0.44	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	15.52	16.61	13.79	-11.14	144	1,044	13.79	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	76.09	74.10	72.70	-4.47	420	559	75.13	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	25.37				153	603	25.37	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.03		0.47	-84.47	4	642	0.62	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Buncombe County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	77.11	84.78	85.68	11.11	2,249	2,625	85.68	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.98	27.38	27.56	-4.91	636	2,308	27.56	28.75
B. Excessive Weight Gain	38.61	40.46	37.74	-2.26	871	2,308	37.74	48.42
C. Inadequate Weight Gain	19.92	20.80	17.59	-11.70	406	2,308	17.59	19.51
D. Unknown Weight Gain	12.49	11.35	17.11	37.04	395	2,308	17.11	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	3.41	4.23	5.18	51.73	136	2,625	5.18	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.86	71.29	72.51	2.34	955	1,357	70.38	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	24.31				301	1,238	24.31	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	9.28		2.01	-78.36	30	1,587	1.89	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Burke County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	64.22	64.29	62.60	-2.53	564	901	62.60	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.53	27.66	29.17	-1.24	231	792	29.17	28.75
B. Excessive Weight Gain	45.05	42.73	37.88	-15.93	300	792	37.88	48.42
C. Inadequate Weight Gain	23.76	24.03	28.41	19.55	225	792	28.41	19.51
D. Unknown Weight Gain	1.65	5.58	4.55	175.76	36	792	4.55	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	19.54	19.89	17.43	-10.81	157	901	17.43	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.15	73.37	77.36	8.72	462	614	75.24	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	31.65				200	632	31.65	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	10.50		9.27	-11.74	18	664	2.71	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Cabarrus County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	61.38	57.34	56.79	-7.46	1,396	2,458	56.79	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.06	27.74	27.84	-7.37	593	2,130	27.84	28.75
B. Excessive Weight Gain	50.03	51.48	49.81	-0.43	1,061	2,130	49.81	48.42
C. Inadequate Weight Gain	18.21	18.76	21.13	15.99	450	2,130	21.13	19.51
D. Unknown Weight Gain	1.71	2.02	1.22	-28.45	26	2,130	1.22	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	9.78	9.29	8.50	-13.02	209	2,458	8.50	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	69.95	66.08	65.65	-6.15	667	1,051	63.46	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	24.13				279	1,156	24.13	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	5.58		8.62	54.50	157	1,249	12.57	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Caldwell County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	68.93	64.25	66.63	-3.35	533	800	66.63	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	24.79	29.83	23.43	-5.48	164	700	23.43	28.75
B. Excessive Weight Gain	52.83	46.59	54.71	3.56	383	700	54.71	48.42
C. Inadequate Weight Gain	22.10	22.73	20.71	-6.25	145	700	20.71	19.51
D. Unknown Weight Gain	0.28	0.85	1.14	303.43	8	700	1.14	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	21.13	23.31	19.38	-8.31	155	800	19.38	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	72.52	79.43	77.28	6.55	446	573	77.84	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	40.51				207	511	40.51	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	24.66		4.54	-81.59	28	600	4.67	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Carteret County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	75.57	72.81	75.98	0.54	465	612	75.98	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.72	30.72	27.97	0.91	146	522	27.97	28.75
B. Excessive Weight Gain	53.26	51.86	55.17	3.59	288	522	55.17	48.42
C. Inadequate Weight Gain	17.93	16.83	16.09	-10.28	84	522	16.09	19.51
D. Unknown Weight Gain	1.09	0.59	0.77	-29.50	4	522	0.77	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.97	17.56	16.67	-12.13	102	612	16.67	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	66.09	71.87	78.57	18.89	238	311	76.53	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	23.74				80	337	23.74	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	15.84		22.90	44.62	90	324	27.78	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Caswell County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	51.85	56.67	63.35	22.18	121	191	63.35	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	24.08	38.55	25.00	3.80	39	156	25.00	28.75
B. Excessive Weight Gain	46.60	41.34	53.21	14.18	83	156	53.21	48.42
C. Inadequate Weight Gain	12.04	17.88	16.67	38.41	26	156	16.67	19.51
D. Unknown Weight Gain	17.28	2.23	5.13	-70.32	8	156	5.13	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.89	20.48	11.52	-17.07	22	191	11.52	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	82.66	69.39	77.21	-6.60	90	124	72.58	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	42.42				56	132	42.42	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						150		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Catawba County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	69.32	70.82	70.75	2.07	1,241	1,754	70.75	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.88	29.37	29.72	-6.77	455	1,531	29.72	28.75
B. Excessive Weight Gain	46.83	51.30	50.42	7.69	772	1,531	50.42	48.42
C. Inadequate Weight Gain	20.34	19.01	19.01	-6.56	291	1,531	19.01	19.51
D. Unknown Weight Gain	0.96	0.32	0.85	-11.15	13	1,531	0.85	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.03	15.84	16.02	-0.05	281	1,754	16.02	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.39	69.91	74.79	1.90	797	1,064	74.91	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	28.40				326	1,148	28.40	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.77		24.83	558.77	315	1,243	25.34	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Chatham County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	52.05	56.39	64.51	23.93	429	665	64.51	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	33.33	29.92	26.89	-19.33	160	595	26.89	28.75
B. Excessive Weight Gain	40.95	42.86	49.41	20.66	294	595	49.41	48.42
C. Inadequate Weight Gain	19.24	23.17	14.29	-25.74	85	595	14.29	19.51
D. Unknown Weight Gain	6.48	4.05	9.41	45.33	56	595	9.41	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	7.72	6.39	3.76	-51.29	25	665	3.76	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.52	70.65	72.38	2.64	153	228	67.11	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	17.46				194	1,111	17.46	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.20		0.90	-24.59	2	345	0.58	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Cherokee County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	64.32	67.62	74.77	16.26	166	222	74.77	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.11	22.16	31.32	11.42	57	182	31.32	28.75
B. Excessive Weight Gain	49.19	58.52	48.35	-1.70	88	182	48.35	48.42
C. Inadequate Weight Gain	20.54	19.32	19.23	-6.38	35	182	19.23	19.51
D. Unknown Weight Gain	2.16	0.00	1.10	-49.18	2	182	1.10	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	22.07	25.24	18.47	-16.30	41	222	18.47	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	81.30	80.34	80.39	-1.12	125	152	82.24	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	40.28				58	144	40.28	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						164		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Clay County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	53.41	75.64	77.11	44.37	64	83	77.11	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	32.47	22.22	33.82	4.18	23	68	33.82	28.75
B. Excessive Weight Gain	45.45	47.62	45.59	0.29	31	68	45.59	48.42
C. Inadequate Weight Gain	15.58	26.98	20.59	32.11	14	68	20.59	19.51
D. Unknown Weight Gain	6.49	3.17	0.00	-100.00	0	68	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	31.82	30.77	25.30	-20.48	21	83	25.30	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	84.21	76.43	90.59	7.57	46	53	86.79	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	25.53				12	47	25.53	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	19.15		17.58	-8.18		57		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Cleveland County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.07	61.14	49.08	-20.93	532	1,084	49.08	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.57	25.91	25.65	0.32	237	924	25.65	28.75
B. Excessive Weight Gain	53.72	51.48	52.49	-2.29	485	924	52.49	48.42
C. Inadequate Weight Gain	20.39	22.17	21.54	5.63	199	924	21.54	19.51
D. Unknown Weight Gain	0.32	0.44	0.32	0.32	3	924	0.32	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	23.76	21.71	21.31	-10.33	231	1,084	21.31	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	83.56	84.82	79.92	-4.36	600	770	77.92	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	22.02				159	722	22.02	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			2.94		25	810	3.09	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Columbus County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	47.39	45.72	38.83	-18.06	233	600	38.83	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.32	26.04	27.75	1.55	144	519	27.75	28.75
B. Excessive Weight Gain	43.17	45.53	43.16	-0.02	224	519	43.16	48.42
C. Inadequate Weight Gain	24.77	23.06	24.47	-1.22	127	519	24.47	19.51
D. Unknown Weight Gain	4.74	5.37	4.62	-2.36	24	519	4.62	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.41	17.27	20.00	21.87	120	600	20.00	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	72.53	76.05	75.95	4.71	344	459	74.95	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	39.43				194	492	39.43	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.05			-100.00		522		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Craven County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	67.02	59.07	62.24	-7.13	905	1,454	62.24	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.25	30.46	28.01	-0.83	358	1,278	28.01	28.75
B. Excessive Weight Gain	50.27	51.22	51.41	2.27	657	1,278	51.41	48.42
C. Inadequate Weight Gain	19.51	17.25	19.56	0.24	250	1,278	19.56	19.51
D. Unknown Weight Gain	1.97	1.07	1.02	-48.47	13	1,278	1.02	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.44	10.87	10.32	-9.79	150	1,454	10.32	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	80.95	84.05	83.64	3.32	509	582	87.46	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	31.48				271	861	31.48	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	12.54		2.22	-82.30	18	619	2.91	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Cumberland County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	74.91	69.37	69.79	-6.83	3,778	5,413	69.79	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.42	28.63	27.56	-3.03	1,264	4,587	27.56	28.75
B. Excessive Weight Gain	53.61	51.90	52.47	-2.12	2,407	4,587	52.47	48.42
C. Inadequate Weight Gain	17.42	19.18	19.69	13.00	903	4,587	19.69	19.51
D. Unknown Weight Gain	0.55	0.29	0.28	-48.55	13	4,587	0.28	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.05	9.51	9.66	-12.58	523	5,413	9.66	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	75.53	74.73	77.64	2.80	1,753	2,261	77.53	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	22.69				824	3,631	22.69	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.47		4.28	191.71	128	2,391	5.35	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Dare County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	63.72	74.36	69.54	9.14	242	348	69.54	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.42	30.29	34.21	8.89	104	304	34.21	28.75
B. Excessive Weight Gain	40.20	42.02	46.38	15.37	141	304	46.38	48.42
C. Inadequate Weight Gain	17.23	22.15	18.42	6.91	56	304	18.42	19.51
D. Unknown Weight Gain	11.15	5.54	0.99	-91.15	3	304	0.99	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.03	10.83	8.91	-11.18	31	348	8.91	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	65.41	64.57	68.35	4.49	104	143	72.73	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	23.89				43	180	23.89	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	20.12		31.63	57.23	66	195	33.85	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Davidson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	73.28	66.57	60.43	-17.52	1,057	1,749	60.43	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.96	29.24	25.43	-12.18	383	1,506	25.43	28.75
B. Excessive Weight Gain	51.29	50.42	49.80	-2.91	750	1,506	49.80	48.42
C. Inadequate Weight Gain	18.09	18.14	21.38	18.18	322	1,506	21.38	19.51
D. Unknown Weight Gain	1.66	2.19	3.39	104.41	51	1,506	3.39	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.55	17.94	15.78	-14.93	276	1,749	15.78	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.83	73.98	75.70	6.88	815	1,082	75.32	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	32.02				350	1,093	32.02	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	13.37		12.95	-3.09	166	1,207	13.75	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Davie County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	80.65	72.58	59.02	-26.82	229	388	59.02	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.73	27.41	28.24	-11.02	96	340	28.24	28.75
B. Excessive Weight Gain	44.55	48.29	48.24	8.27	164	340	48.24	48.42
C. Inadequate Weight Gain	20.51	17.76	15.29	-25.44	52	340	15.29	19.51
D. Unknown Weight Gain	3.21	6.54	8.24	156.94	28	340	8.24	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.35	12.74	9.02	-12.88	35	388	9.02	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.02	70.00	74.32	4.65	153	193	79.27	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	27.06				59	218	27.06	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.56		1.65	5.86	9	228	3.95	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Duplin County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	56.22	59.97	60.29	7.24	451	748	60.29	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.20	30.92	27.70	-1.78	177	639	27.70	28.75
B. Excessive Weight Gain	42.84	46.01	46.95	9.60	300	639	46.95	48.42
C. Inadequate Weight Gain	22.10	22.34	24.88	12.57	159	639	24.88	19.51
D. Unknown Weight Gain	6.86	0.74	0.47	-93.16	3	639	0.47	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	7.63	10.10	8.42	10.38	63	748	8.42	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	72.44	71.47	76.81	6.04	315	408	77.21	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	15.30				88	575	15.30	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	4.66			-100.00		633		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Durham County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	44.25	52.70	64.31	45.33	2,721	4,231	64.31	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.56	29.18	27.58	-3.43	1,006	3,648	27.58	28.75
B. Excessive Weight Gain	45.99	42.82	45.15	-1.82	1,647	3,648	45.15	48.42
C. Inadequate Weight Gain	17.48	19.33	17.98	2.85	656	3,648	17.98	19.51
D. Unknown Weight Gain	7.97	8.67	9.29	16.53	339	3,648	9.29	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	5.22	4.86	4.49	-14.04	190	4,231	4.49	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	62.39	65.26	70.43	12.89	1,060	1,473	71.96	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	19.28				374	1,940	19.28	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	9.47		13.12	38.45	362	2,107	17.18	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Edgecombe County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	50.16	46.92	47.18	-5.94	301	638	47.18	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	22.16	23.90	20.84	-5.94	114	547	20.84	28.75
B. Excessive Weight Gain	48.55	52.99	55.58	14.46	304	547	55.58	48.42
C. Inadequate Weight Gain	26.78	21.91	21.94	-18.09	120	547	21.94	19.51
D. Unknown Weight Gain	2.50	1.20	1.65	-34.31	9	547	1.65	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	17.56	14.98	17.24	-1.83	110	638	17.24	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	84.44	84.98	86.42	2.35	429	508	84.45	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	19.74				91	461	19.74	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						562		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Forsyth County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	83.52	71.75	59.19	-29.13	2,569	4,340	59.19	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.08	26.48	27.29	-6.16	1,012	3,708	27.29	28.75
B. Excessive Weight Gain	46.17	48.13	45.69	-1.04	1,694	3,708	45.69	48.42
C. Inadequate Weight Gain	20.42	16.19	16.91	-17.17	627	3,708	16.91	19.51
D. Unknown Weight Gain	4.33	9.20	10.11	133.32	375	3,708	10.11	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	6.79	5.06	5.41	-20.24	235	4,340	5.41	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	67.02	72.46	70.63	5.39	1,489	2,029	73.39	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	31.64				760	2,402	31.64	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.92			-100.00		2,495		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Franklin County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	48.22	39.91	49.66	2.98	360	725	49.66	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	23.25	25.09	27.70	19.14	172	621	27.70	28.75
B. Excessive Weight Gain	51.28	53.26	47.99	-6.43	298	621	47.99	48.42
C. Inadequate Weight Gain	19.49	16.67	19.97	2.47	124	621	19.97	19.51
D. Unknown Weight Gain	5.98	4.98	4.35	-27.33	27	621	4.35	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.72	12.20	10.21	-12.92	74	725	10.21	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	68.84	65.13	65.67	-4.59	264	392	67.35	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	7.20				28	389	7.20	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	10.50		11.95	13.86	95	446	21.30	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Gaston County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.37	59.30	57.94	-7.10	1,463	2,525	57.94	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.19	26.67	26.98	-0.79	587	2,176	26.98	28.75
B. Excessive Weight Gain	52.69	53.42	51.75	-1.78	1,126	2,176	51.75	48.42
C. Inadequate Weight Gain	18.85	18.46	19.90	5.56	433	2,176	19.90	19.51
D. Unknown Weight Gain	1.27	1.45	1.38	8.35	30	2,176	1.38	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	20.41	20.02	17.82	-12.67	450	2,525	17.82	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	72.78	70.67	70.59	-3.00	1,032	1,469	70.25	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	28.55				438	1,534	28.55	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.79		0.12	-96.84		1,620		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Graham County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	67.78	71.11	74.03	9.22	57	77	74.03	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	24.69	36.00	30.30	22.73	20	66	30.30	28.75
B. Excessive Weight Gain	59.26	42.67	46.97	-20.74	31	66	46.97	48.42
C. Inadequate Weight Gain	12.35	21.33	19.70	59.55	13	66	19.70	19.51
D. Unknown Weight Gain	3.70	0.00	3.03	-18.18	2	66	3.03	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	30.00	34.44	32.47	8.23	25	77	32.47	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	77.01	87.00	89.56	16.29	45	53	84.91	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	16.13				10	62	16.13	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	5.65		2.53	-55.32		57		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Granville-Vance District

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	48.69	50.50	53.17	9.20	596	1,121	53.17	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.68	26.69	23.43	-12.18	224	956	23.43	28.75
B. Excessive Weight Gain	47.17	44.16	48.12	2.00	460	956	48.12	48.42
C. Inadequate Weight Gain	24.01	24.76	24.48	1.93	234	956	24.48	19.51
D. Unknown Weight Gain	2.13	4.39	3.97	86.22	38	956	3.97	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.85	14.49	12.67	-1.43	142	1,121	12.67	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	75.88	72.85	73.77	-2.78	519	736	70.52	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	17.05				125	733	17.05	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	28.21		31.89	13.05	218	813	26.81	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Granville

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	54.37	53.64	61.10	12.38	355	581	61.10	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.67	27.27	24.85	-10.19	124	499	24.85	28.75
B. Excessive Weight Gain	48.15	44.82	50.30	4.47	251	499	50.30	48.42
C. Inadequate Weight Gain	21.35	22.83	20.44	-4.26	102	499	20.44	19.51
D. Unknown Weight Gain	2.83	5.07	4.41	55.67	22	499	4.41	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.55	11.82	10.84	-13.58	63	581	10.84	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	75.85	74.37	74.23	-2.14	227	313	72.52	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	64.18		79.06	23.17	218	341	63.93	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Vance

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	43.52	47.42	44.63	2.54	241	540	44.63	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.73	26.09	21.88	-14.96	100	457	21.88	28.75
B. Excessive Weight Gain	46.23	43.48	45.73	-1.08	209	457	45.73	48.42
C. Inadequate Weight Gain	26.57	26.74	28.88	8.71	132	457	28.88	19.51
D. Unknown Weight Gain	1.46	3.70	3.50	139.07	16	457	3.50	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.13	17.11	14.63	11.45	79	540	14.63	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	75.90	71.78	73.44	-3.24	292	423	69.03	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.48			-100.00		472		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Greene County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	68.29	68.52	63.55	-6.95	129	203	63.55	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.17	27.81	28.73	5.75	52	181	28.73	28.75
B. Excessive Weight Gain	52.02	48.13	46.96	-9.73	85	181	46.96	48.42
C. Inadequate Weight Gain	20.23	22.99	23.20	14.70	42	181	23.20	19.51
D. Unknown Weight Gain	0.58	1.07	1.10	91.16	2	181	1.10	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.66	15.28	13.79	0.99	28	203	13.79	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	82.28	77.64	82.09	-0.22	95	122	77.87	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	35.15				58	165	35.15	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						169		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Guilford County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.23	56.42	58.21	-6.46	3,548	6,095	58.21	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.39	30.69	27.87	-5.18	1,466	5,261	27.87	28.75
B. Excessive Weight Gain	45.74	47.14	47.69	4.28	2,509	5,261	47.69	48.42
C. Inadequate Weight Gain	18.01	17.63	19.86	10.27	1,045	5,261	19.86	19.51
D. Unknown Weight Gain	6.86	4.54	4.58	-33.27	241	5,261	4.58	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	6.78	6.34	5.94	-12.35	362	6,095	5.94	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	63.63	60.57	66.33	4.25	2,224	3,273	67.95	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	27.73				915	3,300	27.73	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	48.06		44.66	-7.09	1,674	3,753	44.60	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Halifax County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	61.11	60.71	60.80	-0.51	349	574	60.80	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.90	24.59	27.94	7.87	133	476	27.94	28.75
B. Excessive Weight Gain	50.53	53.46	50.63	0.20	241	476	50.63	48.42
C. Inadequate Weight Gain	21.44	21.14	20.38	-4.97	97	476	20.38	19.51
D. Unknown Weight Gain	2.12	0.81	1.05	-50.53	5	476	1.05	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	14.52	14.00	13.76	-5.19	79	574	13.76	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	82.46	83.12	83.52	1.29	371	437	84.90	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	34.73				157	452	34.73	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	22.91		17.37	-24.17	51	472	10.81	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Harnett County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	65.95	61.16	60.89	-7.67	1,096	1,800	60.89	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.67	29.39	27.63	-9.92	436	1,578	27.63	28.75
B. Excessive Weight Gain	50.09	50.56	52.47	4.75	828	1,578	52.47	48.42
C. Inadequate Weight Gain	18.37	19.46	18.88	2.82	298	1,578	18.88	19.51
D. Unknown Weight Gain	0.87	0.59	1.01	17.11	16	1,578	1.01	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.26	11.44	9.50	-15.64	171	1,800	9.50	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.65	67.45	72.43	2.53	522	719	72.60	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	23.31				193	828	23.31	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			0.90		24	842	2.85	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Haywood County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	72.48	74.56	75.21	3.77	449	597	75.21	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.06	26.13	22.41	-22.87	117	522	22.41	28.75
B. Excessive Weight Gain	48.50	51.23	45.40	-6.40	237	522	45.40	48.42
C. Inadequate Weight Gain	18.16	18.31	22.41	23.41	117	522	22.41	19.51
D. Unknown Weight Gain	4.27	4.32	9.77	128.62	51	522	9.77	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	20.55	18.51	15.08	-26.64	90	597	15.08	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.01	78.72	76.87	9.80	276	380	72.63	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	41.48				151	364	41.48	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	7.36			-100.00		390		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Henderson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	76.34	75.14	76.24	-0.13	815	1,069	76.24	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.33	30.51	31.05	-0.87	295	950	31.05	28.75
B. Excessive Weight Gain	43.48	45.35	42.53	-2.20	404	950	42.53	48.42
C. Inadequate Weight Gain	19.82	19.96	19.68	-0.71	187	950	19.68	19.51
D. Unknown Weight Gain	5.37	4.18	6.74	25.53	64	950	6.74	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.34	9.82	9.45	-8.67	101	1,069	9.45	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	46.97	51.36	61.70	31.35	356	549	64.85	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.56				170	640	26.56	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	11.72		2.80	-76.15	5	662	0.76	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Hertford County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	65.56	73.54	71.97	9.77	172	239	71.97	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	20.51	26.63	30.00	46.25	57	190	30.00	28.75
B. Excessive Weight Gain	36.92	44.57	33.68	-8.77	64	190	33.68	48.42
C. Inadequate Weight Gain	31.28	26.09	34.74	11.04	66	190	34.74	19.51
D. Unknown Weight Gain	11.28	2.72	1.58	-86.00	3	190	1.58	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	9.96	11.66	13.39	34.45	32	239	13.39	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.45	74.95	84.10	7.21	155	171	90.64	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.18				50	191	26.18	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.60			-100.00		186		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Hoke County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	74.42	69.20	73.12	-1.75	699	956	73.12	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.90	30.19	32.76	13.37	268	818	32.76	28.75
B. Excessive Weight Gain	52.69	49.40	49.88	-5.33	408	818	49.88	48.42
C. Inadequate Weight Gain	17.88	20.05	17.11	-4.26	140	818	17.11	19.51
D. Unknown Weight Gain	0.54	0.36	0.24	-54.52	2	818	0.24	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.12	9.28	9.21	-9.06	88	956	9.21	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	74.44	71.58	75.08	0.85	247	347	71.18	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	8.53				44	516	8.53	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			0.25			404		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Hyde County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	72.41	67.57	79.59	9.91	39	49	79.59	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.27	21.21	26.19	-3.97	11	42	26.19	28.75
B. Excessive Weight Gain	50.00	36.36	54.76	9.52	23	42	54.76	48.42
C. Inadequate Weight Gain	13.64	42.42	19.05	39.68	8	42	19.05	19.51
D. Unknown Weight Gain	9.09	0.00	0.00	-100.00	0	42	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	15.52	5.41	12.24	-21.09	6	49	12.24	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.21	79.12	89.02	25.01	26	28	92.86	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	17.24				5	29	17.24	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			4.90		5	34	14.71	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Iredell County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	61.53	62.96	60.88	-1.06	1,178	1,935	60.88	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.08	29.85	30.23	-2.71	504	1,667	30.23	28.75
B. Excessive Weight Gain	49.61	50.67	48.95	-1.32	816	1,667	48.95	48.42
C. Inadequate Weight Gain	18.59	18.91	20.16	8.40	336	1,667	20.16	19.51
D. Unknown Weight Gain	0.72	0.57	0.66	-8.70	11	1,667	0.66	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.21	12.25	11.21	-8.19	217	1,935	11.21	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	77.34	76.22	71.73	-7.25	676	957	70.64	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	27.62				256	927	27.62	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			0.13		2	1,048	0.19	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Jackson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.26	70.68	65.75	5.61	263	400	65.75	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	24.46	27.41	26.22	7.19	91	347	26.22	28.75
B. Excessive Weight Gain	57.80	53.06	54.18	-6.26	188	347	54.18	48.42
C. Inadequate Weight Gain	15.60	17.49	16.43	5.32	57	347	16.43	19.51
D. Unknown Weight Gain	2.14	2.04	3.17	48.09	11	347	3.17	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	23.97	17.29	19.00	-20.72	76	400	19.00	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	80.20	78.05	74.21	-7.47	168	264	63.64	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	5.24				11	210	5.24	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	0.57		0.11	-80.33		290		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Johnston County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	63.00	62.31	66.39	5.37	1,505	2,267	66.39	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.68	26.84	27.67	-0.01	546	1,973	27.67	28.75
B. Excessive Weight Gain	52.43	53.59	51.39	-1.97	1,014	1,973	51.39	48.42
C. Inadequate Weight Gain	17.75	17.68	17.13	-3.51	338	1,973	17.13	19.51
D. Unknown Weight Gain	2.14	1.89	3.80	77.55	75	1,973	3.80	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	9.61	9.01	8.34	-13.22	189	2,267	8.34	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	54.65	63.92	70.23	28.51	728	1,041	69.93	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	21.88				235	1,074	21.88	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	0.20			-100.00		1,289		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Jones County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	66.67	53.33	65.56	-1.67	59	90	65.56	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	31.87	29.55	32.47	1.88	25	77	32.47	28.75
B. Excessive Weight Gain	43.96	51.14	51.95	18.18	40	77	51.95	48.42
C. Inadequate Weight Gain	21.98	18.18	14.29	-35.00	11	77	14.29	19.51
D. Unknown Weight Gain	2.20	1.14	1.30	-40.91	1	77	1.30	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	15.63	13.33	11.11	-28.89	10	90	11.11	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	79.74	89.60	84.21	5.61	38	46	82.61	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	32.84				22	67	32.84	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	4.98			-100.00		48		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI≥30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Lee County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	49.29	50.95	51.38	4.23	373	726	51.38	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.94	30.11	26.75	-10.65	168	628	26.75	28.75
B. Excessive Weight Gain	45.96	47.62	47.13	2.56	296	628	47.13	48.42
C. Inadequate Weight Gain	22.16	20.12	21.18	-4.41	133	628	21.18	19.51
D. Unknown Weight Gain	1.95	2.15	4.94	153.65	31	628	4.94	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.26	14.87	12.95	-2.33	94	726	12.95	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	66.28	64.42	76.69	15.70	297	381	77.95	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	13.18				63	478	13.18	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.56		6.59	85.12	31	508	6.10	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Lenoir County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	56.15	57.12	59.87	6.64	379	633	59.87	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.98	27.18	24.95	-7.50	134	537	24.95	28.75
B. Excessive Weight Gain	48.63	46.50	49.91	2.64	268	537	49.91	48.42
C. Inadequate Weight Gain	22.68	24.96	23.84	5.10	128	537	23.84	19.51
D. Unknown Weight Gain	1.72	1.37	1.30	-24.13	7	537	1.30	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	15.33	17.62	16.75	9.26	106	633	16.75	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	72.52	72.78	79.66	9.86	363	461	78.74	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.92				140	520	26.92	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.20		0.30	-75.10		518		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Lincoln County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	71.18	68.54	70.42	-1.07	557	791	70.42	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	32.43	26.49	29.28	-9.70	200	683	29.28	28.75
B. Excessive Weight Gain	53.09	55.48	53.73	1.21	367	683	53.73	48.42
C. Inadequate Weight Gain	13.57	17.75	16.84	24.04	115	683	16.84	19.51
D. Unknown Weight Gain	0.90	0.28	0.15	-83.82	1	683	0.15	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.47	13.46	15.93	-3.27	126	791	15.93	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.44	73.66	77.52	5.56	324	416	77.88	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	25.45				113	444	25.45	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	5.05			-100.00		442		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Macon County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	71.98	78.14	71.23	-1.03	260	365	71.23	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.76	30.34	26.47	-7.97	81	306	26.47	28.75
B. Excessive Weight Gain	45.82	47.59	47.06	2.71	144	306	47.06	48.42
C. Inadequate Weight Gain	23.41	21.03	24.51	4.69	75	306	24.51	19.51
D. Unknown Weight Gain	2.01	1.03	1.96	-2.29	6	306	1.96	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	20.06	21.56	18.63	-7.12	68	365	18.63	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	61.94	68.25	79.85	28.91	180	227	79.30	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	17.07				42	246	17.07	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.23			-100.00		269		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Madison County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	80.21	86.73	85.90	7.09	201	234	85.90	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	24.71	27.49	25.13	1.70	50	199	25.13	28.75
B. Excessive Weight Gain	43.53	40.35	40.70	-6.49	81	199	40.70	48.42
C. Inadequate Weight Gain	24.12	24.56	20.60	-14.57	41	199	20.60	19.51
D. Unknown Weight Gain	7.65	7.60	13.57	77.43	27	199	13.57	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	8.02	9.18	11.97	49.17	28	234	11.97	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.09	71.35	77.40	10.42	125	156	80.13	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	34.21				52	152	34.21	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	21.95		4.35	-80.19	1	150	0.67	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Martin-Tyrrell-Washington District

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	68.12	67.10	70.75	3.87	300	424	70.75	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.74	22.22	26.32	-8.43	95	361	26.32	28.75
B. Excessive Weight Gain	50.44	54.05	47.92	-4.99	173	361	47.92	48.42
C. Inadequate Weight Gain	19.65	23.42	25.76	31.12	93	361	25.76	19.51
D. Unknown Weight Gain	1.17	0.30	0.00	-100.00	0	361	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.51	15.68	15.57	-5.74	66	424	15.57	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	86.56	83.24	84.56	-2.30	263	306	85.95	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	15.08				46	305	15.08	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	7.29		7.40	1.47	21	344	6.10	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Martin

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	73.09	69.83	74.44	1.84	198	266	74.44	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.65	20.71	26.43	3.03	60	227	26.43	28.75
B. Excessive Weight Gain	52.88	53.54	50.66	-4.20	115	227	50.66	48.42
C. Inadequate Weight Gain	21.47	25.25	22.91	6.72	52	227	22.91	19.51
D. Unknown Weight Gain	0.00	0.51	0.00		0	227	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.87	18.53	15.79	-6.39	42	266	15.79	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	84.02	82.83	83.40	-0.74	156	185	84.32	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						196		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Tyrrell

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	75.56	65.85	74.36	-1.58	29	39	74.36	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	41.46	18.18	20.00	-51.76	7	35	20.00	28.75
B. Excessive Weight Gain	41.46	66.67	51.43	24.03	18	35	51.43	48.42
C. Inadequate Weight Gain	14.63	15.15	28.57	95.24	10	35	28.57	19.51
D. Unknown Weight Gain	2.44	0.00	0.00	-100.00	0	35	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	22.22	9.76	17.95	-19.23	7	39	17.95	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	90.91	81.94	85.51	-5.94	19	21	90.48	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						29		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Washington

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	57.04	62.07	61.34	7.54	73	119	61.34	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.36	26.47	28.28	-3.66	28	99	28.28	28.75
B. Excessive Weight Gain	49.54	50.98	40.40	-18.44	40	99	40.40	48.42
C. Inadequate Weight Gain	18.35	22.55	31.31	70.66	31	99	31.31	19.51
D. Unknown Weight Gain	2.75	0.00	0.00	-100.00	0	99	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	14.08	12.07	14.29	1.43	17	119	14.29	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	89.60	84.18	86.42	-3.54	88	100	88.00	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	20.82		22.38	7.50	21	119	17.65	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Mecklenburg County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	67.27	62.08	61.73	-8.23	9,168	14,851	61.73	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.30	31.72	31.77	4.85	4,091	12,876	31.77	28.75
B. Excessive Weight Gain	45.62	46.29	46.99	3.00	6,050	12,876	46.99	48.42
C. Inadequate Weight Gain	19.25	19.13	18.70	-2.82	2,408	12,876	18.70	19.51
D. Unknown Weight Gain	4.84	2.86	2.54	-47.49	327	12,876	2.54	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	3.81	3.55	3.34	-12.25	496	14,851	3.34	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	63.95	64.72	59.88	-6.36	3,577	5,621	63.64	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	22.27				1,374	6,171	22.27	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	6.80		9.58	40.92	780	7,280	10.71	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Montgomery County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	58.19	53.77	55.06	-5.38	174	316	55.06	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.97	28.36	26.52	-14.36	74	279	26.52	28.75
B. Excessive Weight Gain	41.04	42.91	44.09	7.41	123	279	44.09	48.42
C. Inadequate Weight Gain	27.61	27.99	28.67	3.85	80	279	28.67	19.51
D. Unknown Weight Gain	0.37	0.75	0.72	92.11	2	279	0.72	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	14.38	16.35	16.46	14.42	52	316	16.46	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.67	77.08	83.66	13.57	174	210	82.86	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	28.16				78	277	28.16	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						266		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Moore County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	60.90	60.22	64.08	5.23	678	1,058	64.08	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.13	31.65	30.89	6.02	282	913	30.89	28.75
B. Excessive Weight Gain	47.98	44.67	51.92	8.21	474	913	51.92	48.42
C. Inadequate Weight Gain	22.43	23.34	16.10	-28.21	147	913	16.10	19.51
D. Unknown Weight Gain	0.46	0.34	1.10	136.86	10	913	1.10	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.32	10.85	11.15	-9.48	118	1,058	11.15	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	50.12	66.80	75.55	50.74	328	442	74.21	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	25.00				124	496	25.00	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	2.11			-100.00		493		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Nash County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	55.53	50.87	51.47	-7.30	541	1,051	51.47	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.27	25.73	25.58	-2.61	231	903	25.58	28.75
B. Excessive Weight Gain	49.12	50.56	51.27	4.38	463	903	51.27	48.42
C. Inadequate Weight Gain	20.89	20.99	22.26	6.56	201	903	22.26	19.51
D. Unknown Weight Gain	3.72	2.71	0.89	-76.20	8	903	0.89	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.76	11.29	12.65	-0.81	133	1,051	12.65	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.07	78.54	83.56	7.02	569	676	84.17	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	23.66				186	786	23.66	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	7.01		1.76	-74.98		768		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
New Hanover County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	67.38	62.64	57.77	-14.27	1,335	2,311	57.77	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.01	28.59	28.08	-6.43	568	2,023	28.08	28.75
B. Excessive Weight Gain	50.49	54.54	52.55	4.07	1,063	2,023	52.55	48.42
C. Inadequate Weight Gain	18.17	16.27	18.54	2.03	375	2,023	18.54	19.51
D. Unknown Weight Gain	1.34	0.60	0.84	-37.20	17	2,023	0.84	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	8.21	8.85	8.61	4.88	199	2,311	8.61	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.41	72.77	72.62	-1.07	726	1,005	72.24	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	23.31				241	1,034	23.31	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	13.89		9.46	-31.90	95	1,179	8.06	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Northampton County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	59.43	63.03	64.41	8.38	114	177	64.41	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.21	26.28	33.77	28.88	51	151	33.77	28.75
B. Excessive Weight Gain	45.52	45.26	44.37	-2.52	67	151	44.37	48.42
C. Inadequate Weight Gain	24.83	25.55	21.19	-14.64	32	151	21.19	19.51
D. Unknown Weight Gain	3.45	2.92	0.66	-80.79	1	151	0.66	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	9.14	15.76	15.82	73.02	28	177	15.82	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	77.07	83.01	86.57	12.33	122	141	86.52	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	24.82				34	137	24.82	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	9.75		4.73	-51.46	11	141	7.80	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Onslow County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	63.54	61.07	69.04	8.65	2,881	4,173	69.04	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.10	30.63	29.31	-2.63	1,095	3,736	29.31	28.75
B. Excessive Weight Gain	54.09	53.97	54.42	0.60	2,033	3,736	54.42	48.42
C. Inadequate Weight Gain	15.35	15.15	15.63	1.87	584	3,736	15.63	19.51
D. Unknown Weight Gain	0.46	0.25	0.64	39.54	24	3,736	0.64	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	7.53	6.87	6.61	-12.19	276	4,173	6.61	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	74.84	76.17	77.28	3.26	832	1,092	76.19	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	17.97				432	2,404	17.97	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	20.11		27.95	38.99	324	1,140	28.42	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Orange County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	57.72	57.45	64.95	12.54	784	1,207	64.95	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	32.90	33.55	32.39	-1.55	342	1,056	32.39	28.75
B. Excessive Weight Gain	40.65	39.87	41.10	1.09	434	1,056	41.10	48.42
C. Inadequate Weight Gain	20.00	22.03	17.05	-14.77	180	1,056	17.05	19.51
D. Unknown Weight Gain	6.45	4.55	9.47	46.85	100	1,056	9.47	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	5.54	4.20	3.31	-40.15	40	1,207	3.31	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	68.75	67.85	71.17	3.52	263	374	70.32	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	17.46				194	1,111	17.46	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	15.77		10.67	-32.36	54	485	11.13	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Pamlico County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.03	61.54	52.38	-15.55	44	84	52.38	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.09	17.57	24.66	-5.48	18	73	24.66	28.75
B. Excessive Weight Gain	37.68	55.41	52.05	38.15	38	73	52.05	48.42
C. Inadequate Weight Gain	34.78	25.68	21.92	-36.99	16	73	21.92	19.51
D. Unknown Weight Gain	1.45	1.35	1.37	-5.48	1	73	1.37	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	17.72	13.19	25.00	41.07	21	84	25.00	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	90.36	81.77	89.94	-0.46	52	56	92.86	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	34.29				24	70	34.29	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	2.45			-100.00		75		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Pender County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	58.41	55.84	54.63	-6.46	336	615	54.63	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.35	30.04	29.79	8.94	157	527	29.79	28.75
B. Excessive Weight Gain	51.43	49.29	51.23	-0.38	270	527	51.23	48.42
C. Inadequate Weight Gain	18.37	19.61	18.03	-1.86	95	527	18.03	19.51
D. Unknown Weight Gain	2.86	1.06	0.95	-66.79	5	527	0.95	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.23	13.09	11.71	14.49	72	615	11.71	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	70.00	65.85	73.58	5.11	227	301	75.42	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	31.45				106	337	31.45	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	4.06		14.95	268.62	36	366	9.84	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Person County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	53.26	51.49	66.00	23.94	266	403	66.00	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.00	21.58	29.51	18.05	103	349	29.51	28.75
B. Excessive Weight Gain	45.83	51.58	44.70	-2.47	156	349	44.70	48.42
C. Inadequate Weight Gain	24.22	21.84	21.78	-10.08	76	349	21.78	19.51
D. Unknown Weight Gain	4.95	5.00	4.01	-18.93	14	349	4.01	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.14	14.48	14.14	-22.03	57	403	14.14	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	83.60	83.61	81.72	-2.26	206	251	82.07	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	33.47				84	251	33.47	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	61.54		49.58	-19.43	67	265	25.28	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Pitt County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	74.23	75.25	75.13	1.21	1,625	2,163	75.13	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.92	28.70	29.12	4.30	530	1,820	29.12	28.75
B. Excessive Weight Gain	52.59	52.17	50.38	-4.20	917	1,820	50.38	48.42
C. Inadequate Weight Gain	18.69	18.63	20.11	7.60	366	1,820	20.11	19.51
D. Unknown Weight Gain	0.80	0.50	0.38	-51.79	7	1,820	0.38	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.61	10.16	10.12	-12.76	219	2,163	10.12	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	75.50	75.02	74.12	-1.83	917	1,210	75.79	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	18.82				224	1,190	18.82	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	31.42		29.33	-6.65	463	1,307	35.42	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Randolph County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	53.49	52.78	55.26	3.31	861	1,558	55.26	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.28	28.38	27.68	1.47	377	1,362	27.68	28.75
B. Excessive Weight Gain	48.76	50.53	50.73	4.04	691	1,362	50.73	48.42
C. Inadequate Weight Gain	19.29	18.75	18.65	-3.34	254	1,362	18.65	19.51
D. Unknown Weight Gain	4.66	2.34	2.94	-37.04	40	1,362	2.94	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	14.79	14.72	14.18	-4.09	221	1,558	14.18	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	69.13	73.21	72.78	5.28	684	925	73.95	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	36.23				342	944	36.23	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	6.27			-100.00		1,061		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Richmond County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	58.40	59.52	51.71	-11.46	288	557	51.71	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	27.49	26.51	23.40	-14.88	110	470	23.40	28.75
B. Excessive Weight Gain	49.22	46.05	50.43	2.44	237	470	50.43	48.42
C. Inadequate Weight Gain	23.06	27.21	25.11	8.87	118	470	25.11	19.51
D. Unknown Weight Gain	0.22	0.23	1.06	379.79	5	470	1.06	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	24.25	23.81	24.42	0.67	136	557	24.42	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	77.68	79.28	84.50	8.78	367	441	83.22	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	22.44				92	410	22.44	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	17.60		1.21	-93.15		488		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Robeson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	54.45	54.92	46.30	-14.96	820	1,771	46.30	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.46	29.04	28.83	-2.15	431	1,495	28.83	28.75
B. Excessive Weight Gain	41.20	38.44	40.54	-1.61	606	1,495	40.54	48.42
C. Inadequate Weight Gain	28.23	30.95	29.70	5.21	444	1,495	29.70	19.51
D. Unknown Weight Gain	1.11	1.58	0.94	-15.77	14	1,495	0.94	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	19.54	18.64	17.73	-9.29	314	1,771	17.73	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	75.06	72.20	74.08	-1.31	1,028	1,363	75.42	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	30.01				401	1,336	30.01	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	7.01		4.01	-42.82	103	1,541	6.68	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Rockingham County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	78.91	68.43	72.93	-7.58	660	905	72.93	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.06	28.54	25.16	0.39	198	787	25.16	28.75
B. Excessive Weight Gain	51.93	50.77	51.97	0.08	409	787	51.97	48.42
C. Inadequate Weight Gain	18.07	18.67	20.84	15.31	164	787	20.84	19.51
D. Unknown Weight Gain	4.94	2.02	2.03	-58.84	16	787	2.03	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	17.21	14.95	14.70	-14.60	133	905	14.70	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.63	78.34	78.00	-0.79	450	597	75.38	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	43.23				249	576	43.23	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	31.79		28.66	-9.83	147	649	22.65	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Rowan County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	55.04	52.55	54.21	-1.50	869	1,603	54.21	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.01	28.02	28.77	2.71	397	1,380	28.77	28.75
B. Excessive Weight Gain	51.26	50.33	49.28	-3.88	680	1,380	49.28	48.42
C. Inadequate Weight Gain	19.69	20.48	21.01	6.74	290	1,380	21.01	19.51
D. Unknown Weight Gain	1.04	1.17	0.94	-9.43	13	1,380	0.94	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.04	17.09	15.66	-13.19	251	1,603	15.66	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	69.24	68.94	67.86	-1.99	662	961	68.89	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	35.24				321	911	35.24	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						1,063		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Rutherford-Polk-McDowell District

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	71.61	71.76	68.85	-3.84	902	1,310	68.85	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.34	28.70	27.00	-7.96	310	1,148	27.00	28.75
B. Excessive Weight Gain	48.75	46.48	49.74	2.03	571	1,148	49.74	48.42
C. Inadequate Weight Gain	20.93	23.29	20.99	0.30	241	1,148	20.99	19.51
D. Unknown Weight Gain	0.98	1.53	2.26	130.19	26	1,148	2.26	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	22.67	21.25	20.00	-11.76	262	1,310	20.00	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.72	78.92	81.80	14.05	751	925	81.19	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	24.20				218	901	24.20	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	36.51		38.42	5.25	322	975	33.03	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Rutherford

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	71.49	69.05	63.29	-11.48	431	681	63.29	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.21	26.88	25.33	-10.20	152	600	25.33	28.75
B. Excessive Weight Gain	48.14	49.58	55.17	14.59	331	600	55.17	48.42
C. Inadequate Weight Gain	23.48	22.87	18.50	-21.21	111	600	18.50	19.51
D. Unknown Weight Gain	0.17	0.67	1.00	492.00	6	600	1.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	23.53	21.58	21.73	-7.64	148	681	21.73	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.14	83.66	83.91	17.95	440	517	85.11	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	64.03		71.08	11.00	322	521	61.80	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Polk

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	72.19	60.26	56.43	-21.83	79	140	56.43	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	35.29	33.09	26.02	-26.29	32	123	26.02	28.75
B. Excessive Weight Gain	45.38	47.06	52.03	14.66	64	123	52.03	48.42
C. Inadequate Weight Gain	17.65	16.91	19.51	10.57	24	123	19.51	19.51
D. Unknown Weight Gain	1.68	2.94	2.44	45.12	3	123	2.44	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	15.89	14.74	15.71	-1.13	22	140	15.71	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	64.50	67.58	73.40	13.80	64	86	74.42	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.82			-100.00		86		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

McDowell

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	71.58	79.87	80.16	11.99	392	489	80.16	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.24	30.03	29.65	1.40	126	425	29.65	28.75
B. Excessive Weight Gain	50.61	41.29	41.41	-18.18	176	425	41.41	48.42
C. Inadequate Weight Gain	18.18	26.27	24.94	37.18	106	425	24.94	19.51
D. Unknown Weight Gain	1.97	2.41	4.00	103.50	17	425	4.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	23.64	23.04	18.81	-20.43	92	489	18.81	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	74.62	74.15	80.95	8.49	247	322	76.71	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.58			-100.00		368		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Sampson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	64.75	66.90	67.55	4.31	562	832	67.55	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.86	27.46	27.06	-9.38	200	739	27.06	28.75
B. Excessive Weight Gain	45.68	47.61	49.53	8.43	366	739	49.53	48.42
C. Inadequate Weight Gain	22.97	23.24	21.92	-4.58	162	739	21.92	19.51
D. Unknown Weight Gain	1.49	1.69	1.49	0.14	11	739	1.49	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.53	11.67	12.02	-4.07	100	832	12.02	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	78.68	71.92	76.16	-3.21	384	469	81.88	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	5.46				29	531	5.46	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	8.62		8.88	3.04	77	658	11.70	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Scotland County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	75.89	68.18	58.01	-23.56	257	443	58.01	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.91	34.70	14.05	-47.78	52	370	14.05	28.75
B. Excessive Weight Gain	42.74	40.98	70.00	63.77	259	370	70.00	48.42
C. Inadequate Weight Gain	29.82	23.77	15.68	-47.42	58	370	15.68	19.51
D. Unknown Weight Gain	0.53	0.55	0.27	-48.78	1	370	0.27	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	21.43	21.82	18.06	-15.73	80	443	18.06	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	84.25	84.52	86.09	2.18	316	368	85.87	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	20.52				87	424	20.52	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	15.98			-100.00		385		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Stanly County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	69.75	64.07	66.37	-4.84	450	678	66.37	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.43	30.00	34.26	12.60	197	575	34.26	28.75
B. Excessive Weight Gain	51.60	50.16	47.65	-7.65	274	575	47.65	48.42
C. Inadequate Weight Gain	17.79	19.68	18.09	1.65	104	575	18.09	19.51
D. Unknown Weight Gain	0.18	0.16	0.00	-100.00	0	575	0.00	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	12.07	13.79	16.08	33.21	109	678	16.08	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	81.71	76.71	74.92	-8.31	316	425	74.35	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	34.55				152	440	34.55	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	2.40		1.24	-48.26	16	457	3.50	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Stokes County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	87.20	73.38	63.50	-27.17	261	411	63.50	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.05	19.67	26.51	-8.74	92	347	26.51	28.75
B. Excessive Weight Gain	48.93	55.74	44.38	-9.30	154	347	44.38	48.42
C. Inadequate Weight Gain	19.27	16.67	19.60	1.72	68	347	19.60	19.51
D. Unknown Weight Gain	2.75	7.92	9.51	245.53	33	347	9.51	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	19.47	14.18	14.11	-27.51	58	411	14.11	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	76.00	69.33	72.71	-4.32	163	241	67.63	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	36.44				86	236	36.44	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)			1.16			255		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Surry County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	79.39	71.11	68.00	-14.35	510	750	68.00	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.74	30.41	29.61	10.72	188	635	29.61	28.75
B. Excessive Weight Gain	49.53	48.39	48.35	-2.38	307	635	48.35	48.42
C. Inadequate Weight Gain	22.15	18.71	18.27	-17.53	116	635	18.27	19.51
D. Unknown Weight Gain	1.58	2.49	3.78	138.87	24	635	3.78	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	21.44	21.37	25.33	18.17	190	750	25.33	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	68.41	70.69	73.40	7.31	377	510	73.92	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	25.25				124	491	25.25	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	51.40		31.88	-37.98	168	575	29.22	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Swain County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	55.83	61.00	56.16	0.61	123	219	56.16	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.11	24.55	28.65	1.92	53	185	28.65	28.75
B. Excessive Weight Gain	54.59	58.08	44.32	-18.81	82	185	44.32	48.42
C. Inadequate Weight Gain	16.22	16.17	23.78	46.67	44	185	23.78	19.51
D. Unknown Weight Gain	1.08	1.20	3.24	200.00	6	185	3.24	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	26.70	29.00	25.11	-5.94	55	219	25.11	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	85.38	77.88	66.13	-22.55	83	169	49.11	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	24.69				20	81	24.69	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	0.62		2.15	248.76	2	170	1.18	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Toe River District

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	69.91	71.56	74.38	6.38	357	480	74.38	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	26.46	29.18	26.03	-1.62	107	411	26.03	28.75
B. Excessive Weight Gain	45.29	45.62	47.20	4.22	194	411	47.20	48.42
C. Inadequate Weight Gain	25.45	21.22	21.17	-16.81	87	411	21.17	19.51
D. Unknown Weight Gain	2.80	3.98	5.60	99.93	23	411	5.60	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.81	20.75	19.38	15.23	93	480	19.38	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	72.00	73.54	78.57	9.13	220	283	77.74	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	32.35				99	306	32.35	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.65		0.62	-62.38		323		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Avery

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	75.66	73.68	79.59	5.20	117	147	79.59	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	34.62	23.23	23.20	-32.98	29	125	23.20	28.75
B. Excessive Weight Gain	45.38	49.49	52.80	16.34	66	125	52.80	48.42
C. Inadequate Weight Gain	18.46	23.23	22.40	21.33	28	125	22.40	19.51
D. Unknown Weight Gain	1.54	4.04	1.60	4.00	2	125	1.60	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.82	16.67	14.97	8.33	22	147	14.97	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.68	72.76	86.29	17.11	72	83	86.75	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	3.28		1.67	-49.00		102		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Mitchell

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	64.79	64.79	72.85	12.44	110	151	72.85	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	22.76	33.61	28.91	26.98	37	128	28.91	28.75
B. Excessive Weight Gain	42.28	49.18	46.09	9.03	59	128	46.09	48.42
C. Inadequate Weight Gain	29.27	13.93	21.88	-25.26	28	128	21.88	19.51
D. Unknown Weight Gain	5.69	3.28	3.13	-45.09	4	128	3.13	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	16.20	26.76	25.83	59.46	39	151	25.83	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	74.09	75.93	74.73	0.86	64	93	68.82	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.07			-100.00		99		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

**Process Outcome Objectives for Counties Composing Districts
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA**

Yancey

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	68.99	75.72	71.43	3.54	130	182	71.43	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	22.14	29.49	25.95	17.19	41	158	25.95	28.75
B. Excessive Weight Gain	47.86	40.38	43.67	-8.75	69	158	43.67	48.42
C. Inadequate Weight Gain	28.57	25.64	19.62	-31.33	31	158	19.62	19.51
D. Unknown Weight Gain	1.43	4.49	10.76	653.16	17	158	10.76	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	20.25	18.50	17.58	-13.19	32	182	17.58	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	68.64	72.42	75.98	10.69	84	107	78.50	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]								25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	0.78		0.26	-66.05		122		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Transylvania County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	77.54	76.82	79.93	3.07	219	274	79.93	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	33.20	24.80	30.25	-8.88	72	238	30.25	28.75
B. Excessive Weight Gain	44.40	49.59	42.86	-3.47	102	238	42.86	48.42
C. Inadequate Weight Gain	20.40	21.95	15.55	-23.79	37	238	15.55	19.51
D. Unknown Weight Gain	2.00	3.66	11.34	467.23	27	238	11.34	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	13.33	15.57	9.49	-28.83	26	274	9.49	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.43	68.14	71.79	0.50	131	183	71.58	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	30.73				55	179	30.73	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	4.96		3.86	-22.28	5	203	2.46	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Union County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	64.74	67.24	66.60	2.86	1,571	2,359	66.60	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	30.53	30.55	31.15	2.01	641	2,058	31.15	28.75
B. Excessive Weight Gain	50.35	50.07	49.27	-2.13	1,014	2,058	49.27	48.42
C. Inadequate Weight Gain	18.38	18.66	19.19	4.43	395	2,058	19.19	19.51
D. Unknown Weight Gain	0.74	0.72	0.39	-47.55	8	2,058	0.39	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	7.65	6.77	7.12	-6.85	168	2,359	7.12	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	54.76	59.46	60.63	10.71	492	857	57.41	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	30.04				304	1,012	30.04	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	8.58		7.88	-8.11	88	1,108	7.94	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Wake County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	55.49	54.47	60.78	9.55	7,821	12,867	60.78	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	32.13	30.99	30.97	-3.61	3,494	11,282	30.97	28.75
B. Excessive Weight Gain	45.50	48.34	46.89	3.05	5,290	11,282	46.89	48.42
C. Inadequate Weight Gain	18.71	17.54	17.47	-6.61	1,971	11,282	17.47	19.51
D. Unknown Weight Gain	3.66	3.13	4.67	27.47	527	11,282	4.67	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	3.18	2.79	2.75	-13.58	354	12,867	2.75	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	68.67	68.98	67.82	-1.24	2,554	3,675	69.50	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.18				1,089	4,160	26.18	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	21.46		27.62	28.75	1,355	4,716	28.73	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Warren County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	44.09	44.44	56.99	29.27	106	186	56.99	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	25.63	24.68	21.52	-16.02	34	158	21.52	28.75
B. Excessive Weight Gain	51.25	44.30	50.63	-1.20	80	158	50.63	48.42
C. Inadequate Weight Gain	18.13	25.32	25.95	43.17	41	158	25.95	19.51
D. Unknown Weight Gain	5.00	5.70	1.90	-62.03	3	158	1.90	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	15.05	16.16	12.90	-14.29	24	186	12.90	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	85.78	81.39	82.49	-3.83	131	148	88.51	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	26.35				39	148	26.35	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	1.10		1.41	28.40		157		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Wayne County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	62.38	59.93	56.75	-9.03	925	1,630	56.75	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.80	31.11	29.63	2.88	416	1,404	29.63	28.75
B. Excessive Weight Gain	48.64	46.93	44.66	-8.19	627	1,404	44.66	48.42
C. Inadequate Weight Gain	21.83	21.75	25.21	15.49	354	1,404	25.21	19.51
D. Unknown Weight Gain	0.73	0.20	0.50	-31.70	7	1,404	0.50	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.15	9.73	10.43	2.73	170	1,630	10.43	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	77.03	73.03	74.12	-3.77	675	912	74.01	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	10.17				116	1,141	10.17	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	22.41		23.50	4.85	250	1,082	23.11	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Wilkes County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	75.04	69.53	72.46	-3.43	521	719	72.46	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	33.15	27.35	29.30	-11.63	179	611	29.30	28.75
B. Excessive Weight Gain	44.69	46.62	46.81	4.74	286	611	46.81	48.42
C. Inadequate Weight Gain	21.43	25.37	23.08	7.69	141	611	23.08	19.51
D. Unknown Weight Gain	0.73	0.66	0.82	11.70	5	611	0.82	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.91	19.68	19.47	2.94	140	719	19.47	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	81.06	77.73	79.65	-1.74	383	477	80.29	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	33.19				154	464	33.19	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	31.75		14.37	-54.73	102	521	19.58	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Wilson County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	60.98	55.13	57.05	-6.44	554	971	57.05	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	28.04	27.81	28.05	0.03	230	820	28.05	28.75
B. Excessive Weight Gain	47.89	47.74	41.71	-12.91	342	820	41.71	48.42
C. Inadequate Weight Gain	23.05	23.67	29.51	28.05	242	820	29.51	19.51
D. Unknown Weight Gain	1.02	0.78	0.73	-28.57	6	820	0.73	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	11.19	10.68	8.86	-20.88	86	971	8.86	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	84.30	83.62	80.43	-4.59	477	606	78.71	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	15.42				101	655	15.42	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	5.60		2.10	-62.50		724		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
Yadkin County

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data			NC Percent or Rate
	1st	2nd	3rd		Num	Den	Percent or Rate	
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	75.94	68.33	59.07	-22.22	228	386	59.07	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)								
A. Adequate Weight Gain	29.36	27.87	27.27	-7.11	93	341	27.27	28.75
B. Excessive Weight Gain	44.19	48.52	49.56	12.16	169	341	49.56	48.42
C. Inadequate Weight Gain	22.09	19.02	17.89	-19.03	61	341	17.89	19.51
D. Unknown Weight Gain	4.36	4.59	5.28	21.06	18	341	5.28	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	18.30	16.42	12.95	-29.20	50	386	12.95	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	73.81	77.15	73.57	-0.32	159	216	73.61	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	30.88				67	217	30.88	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)						254		12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.

Process Outcome Objectives
SFY2017-2018 MATERNAL HEALTH AGREEMENT ADDENDA DATA
North Carolina

Process Outcome Objectives	Time Period			% Change from 1st period to 3rd period	Most Recent Data		
	1st	2nd	3rd		Num	Den	Percent or Rate
1. Percentage of women having live births who had adequate prenatal care as defined by Kessner Index. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	64.17	62.04	62.89	-2.00	75,984	120,826	62.89
2. Percentage of women with live term singleton births who gained adequate/excessive/ inadequate weight according to the National Academy of Sciences-Institute of Medicine (IOM) 2009 Recommended Total Weight Gain Ranges during Pregnancy.[2] (Time Periods: 1st - CY13 2nd - CY14; 3rd - CY15. Most Recent - CY15)							
A. Adequate Weight Gain	29.24	29.26	28.75	-1.70	30,040	104,501	28.75
B. Excessive Weight Gain	47.91	48.53	48.42	1.05	50,597	104,501	48.42
C. Inadequate Weight Gain	19.37	19.42	19.51	0.75	20,392	104,501	19.51
D. Unknown Weight Gain	3.48	2.79	3.32	-4.41	3,472	104,501	3.32
3. Percentage of women having live births who smoked during pregnancy. [1] (Time Periods: 1st - CY13; 2nd - CY14; 3rd - CY15. Most Recent - CY15)	10.29	9.84	9.35	-9.13	11,297	120,826	9.35
4. Percentage of Medicaid enrolled pregnant women who deliver who receive prenatal WIC Program services.[3] (Time Periods: 1st - CY07-09; 2nd - CY10-12; 3rd - CY13-15. Most Recent - CY15)	71.39	71.53	72.76	1.91	42,516	58,086	73.19
5. Percentage of infants enrolled in WIC who breastfed at 6 weeks postpartum. (Time Periods: 1st - FY15; 2nd - N/A; 3rd - N/A. Most Recent - FY15) [4]	25.14				16,449	65,434	25.14
6. Percentage of Medicaid enrolled pregnant women who deliver who receive a postpartum home visit.[3] (Time Periods: 1st - CY10-12; 2nd - N/A; 3rd - CY13-15. Most Recent - CY15)	12.19		11.79	-3.31	8,315	68,077	12.21

[1] With the implementation of the 2003 revised US Standard Certificate of Live Birth in NC in August 2010, data on prenatal care initiation and tobacco use prior to 2011 are not comparable.

[2] Recommended total prenatal weight gain range by prepregnancy weight category is 28-40 lbs. for underweight women (BMI<18.5); 25-35 lbs. for normal weight women (BMI 18.5 to 24.9); 15-25 lbs. for overweight women (BMI 25 to 29.9); and 11-20 lbs. for obese women (BMI>=30).

[3] These data exclude women enrolled in Medicaid only for emergency delivery. 2014 Medicaid match includes data from both legacy WIC and Crossroads WIC System. Effective 2015 Prenatal Medicaid match included only Crossroads WIC data.

[4] Due to Crossroads implementation, data are only available for this time period. This page will be updated to include FY16 data when it becomes available. Also, data for counties composing districts are not available.

Data sources for these objectives are as follows: 1, 2, & 3 NC Birth Certificates; 4 NC Automated WIC System matched with Medicaid Processed Claims Report; 5 NC Automated WIC System; and 6 Medicaid Processed Claims Report.